

Practice Parameter for the Assessment and Treatment of Children and Adolescents With Reactive Attachment Disorder and Disinhibited Social Engagement Disorder

Charles H. Zeanah, MD, Tessa Chesher, DO, Neil W. Boris, MD, and the American Academy of Child and Adolescent Psychiatry (AACAP) Committee on Quality Issues (CQI)

This Practice Parameter is a revision of a previous Parameter addressing reactive attachment disorder that was published in 2005. It reviews the current status of reactive attachment disorder (RAD) and disinhibited social engagement disorder (DSED) with regard to assessment and treatment. Attachment is a central component of social and emotional development in early childhood, and disordered attachment is defined by specific patterns of abnormal social behavior in the context of “insufficient care” or social neglect. Assessment requires direct observation of the child in the

context of his or her relationships with primary caregivers. Treatment requires establishing an attachment relationship for the child when none exists and ameliorating disturbed social relatedness with non-caregivers when evident.

Key words: reactive attachment disorder, disinhibited social engagement disorder, indiscriminate behavior, indiscriminate friendliness, indiscriminate social behavior

J Am Acad Child Adolesc Psychiatry 2016;55(11):990–1003.

The original AACAP Parameter on reactive attachment disorder (RAD) was published in 2005.¹ Since then, considerable research has been conducted, and the *DSM-5*² presents what used to be designated as RAD as two distinct disorders—reactive attachment disorder and disinhibited social engagement disorder (DSED). This Parameter updates the 2005 version to incorporate new findings relevant to both disorders.

The majority of research on RAD and DSED has been conducted on children between the ages of 1 and 5 years old. There is little research about homotypic continuity of these disorders or even a consensus about how they manifest in older children and adolescents. Hence, this Parameter focuses primarily on preschool children, with some discussion of the disorders in older children.

Reports linking extremely adverse caregiving environments with aberrant social behaviors in young children date back more than a century. Defining disorders of attachment in formal nosologies began with the *DSM-III* in 1980.³ The *DSM-5*² has defined DSED and RAD as two separate and distinct disorders grouped with other trauma-related disorders. Together, RAD and DSED capture distinctive patterns of aberrant attachment and social behaviors in young children who are socially neglected or are being raised in environments that limit opportunities to form selective attachments. The empirical basis for these disorders is growing, and data are informing a growing consensus about principles of assessment and safe and effective interventions for RAD and DSED.

METHODOLOGY

The initial literature search was conducted in May 2012 using PubMed, PsycINFO, and Cochrane Library databases. The initial search was limited to the date range January 1, 1998, to May 17, 2012. The search in PubMed used the MeSH term “reactive attachment disorders,” which yielded 307 results, and the following key words linked with the Boolean “OR”: “Reactive Attachment Disorders,” “Attachment Disorders,” “Indiscriminate Behavior,” “Indiscriminate Friendliness,” “Indiscriminate Socially Disinhibited Reactive Attachment Disorder,” and “Disinhibited Social Engagement Disorder,” which yielded 361 results. The search in PsycINFO used the thesaurus term “attachment disorder” and yielded 393 results. An advanced search linking keywords with the Boolean “OR” using the terms “Reactive Attachment Disorder,” “Attachment Disorders,” “Indiscriminate Behavior,” “Indiscriminate Friendliness,” “Indiscriminate Socially Disinhibited Reactive Attachment Disorder,” and “Disinhibited Social Engagement Disorder” yielded 287 results. The search was repeated in the Cochrane Library Database, yielding 2 results.

The initial search was extended in June 2016 using PubMed, PsycINFO, and Cochrane Library databases to cover the date range May 18, 2012, to June 15, 2016. The search in PubMed used the MeSH term “reactive attachment disorders,” which yielded 110 results, and the following key words linked with the Boolean “OR”: “Reactive Attachment Disorders,” “Attachment Disorders,” “Indiscriminate Behavior,” “Indiscriminate Friendliness,” “Indiscriminate Socially Disinhibited Reactive Attachment Disorder,” and “Disinhibited Social Engagement Disorder,” which yielded 154 results. The search in PsycINFO used the thesaurus term “attachment disorder” and yielded 119 results. An advanced search linking keywords with the Boolean “OR” using the terms



This article can be used to obtain continuing medical education (CME) at www.jaacap.org.

“Reactive Attachment Disorder,” “Attachment Disorders,” “Indiscriminate Behavior,” “Indiscriminate Friendliness,” “Indiscriminate Socially Disinhibited Reactive Attachment Disorder,” and “Disinhibited Social Engagement Disorder” yielded 187 results. The search was repeated in the Cochrane Library Database of Systematic Reviews, yielding 0 results (when the additional keywords were added and linked with the Boolean “OR,” the search yielded 2 results).

Reviewers of the titles and abstracts of all articles examined key quality domains, including descriptions of the study population (inclusion and exclusion criteria), randomization, blinding, interventions, outcomes (including “last observation carried forward” data and description of dropouts), and statistical analysis. For this Practice Parameter, 83 publications from the first search and 40 from the second search were selected for careful examination on the basis of their weight in the hierarchy of evidence based on quality and relevance to clinical practice.

In addition, searches of relevant publications by the following authors were conducted because of their expertise in this area: Neil W. Boris, Kim Chisholm, Patricia Crittenden, Mary Dozier, Marinus van IJzendoorn, Alicia Lieberman, Karlen Lyons-Ruth, Mary Margaret Gleason, Helen Minnis, Thomas O’Connor, Michael Rutter, Anna Smyke, Isabel Soares, and Charles H. Zeanah. The titles and abstracts of all articles were reviewed.

ATTACHMENT AND ITS DEVELOPMENT

Attachment between an infant and his or her primary caregivers is a biologically driven process that results in organization of behaviors in the young child, especially behavior designed to achieve physical proximity to a preferred caregiver when the child is in need of comfort, support, nurturance, or protection.⁴ The process of attachment unfolds in the first years of life. Newborns recognize their mother’s smell and sound soon after birth but express no preference for a particular person to provide comfort for distress. Between 2 and 7 months of age, infants are motivated to interact socially with a variety of partners, familiar and unfamiliar. During this time, the infant may be more readily comforted by a familiar caregiver, although he or she is generally able to be soothed by unfamiliar adults as well. However, at around 7 to 9 months, infants begin to exhibit reticence around unfamiliar adults (stranger wariness) and to protest separations from familiar caregivers (separation protest). Once these behaviors have appeared, the infant is said to have formed a selective or preferred attachment.

Infants become attached to caregivers with whom they have had significant amounts of interaction.⁵ Although no definitive data are available, this appears to be a relatively small number of adults whom the infant learns through experience that he or she can count on to provide comfort, support, nurturance, and protection, especially in times of stress. These attachment figures appear to be arranged hierarchically by the infant in terms of strength of preference, so that the infant has a most preferred caregiver, a next most preferred caregiver, etc.⁶ There appear to be limits to infants’ capacities to attach to large numbers of caregivers,

presumably because actual physical contact and regular interaction seems to be required for attachments to form. In institutions in which large numbers of caregivers work irregular shifts, many children develop little or no preference for one or more attachment figures.⁷⁻¹⁰ Nevertheless, the limit of how many different attachment figures an infant can have without problems ensuing is unknown.

Preferred attachments to caregivers may develop at any time after infants reach a developmental age of 7 to 9 months, provided that the caregivers have sufficient involvement with the child. Thus, young children removed from institutions or from neglecting families readily form attachments to their new caregivers,^{9,11-15} although the quality of these subsequent attachments can be compromised.^{12,13}

By 12 months of age, it becomes possible to assess the quality of an infant’s attachment to a preferred attachment figure. A laboratory paradigm, the Strange Situation Procedure,¹⁶ involves a series of interactions between a young child, an attachment figure, and an unfamiliar adult, including separations and reunions. During this procedure, individual differences in the organization of 11- to 20-month-old infants’ attachment behaviors as they are directed toward an attachment figure are described in four patterns of attachment—secure, avoidant, resistant, and disorganized.

The Strange Situation Procedure has been conducted in many cultures throughout the world. Although there is variability in distributions within and across different cultures, the same four patterns are evident.¹⁷ These patterns of attachment are relationship-specific rather than within-the-child traits in that the same child’s pattern of attachment may be different with different caregiving adults.¹⁸ These patterns have been associated with different types of caregiving in the first year of life¹⁹ and with differing adaptation in the preschool years and beyond.^{19,20}

For children 2 to 4½ years old, the MacArthur system (J. Cassidy, R. Marvin, unpublished, 1992) describes secure, avoidant, dependent (ambivalent), controlling, and insecure/other patterns of attachment. These classifications are derived from the same Strange Situation Procedure administered to infants, but the classifications are defined by different behaviors, with increased reliance on verbal rather than only motor behaviors.

Insecure attachment (avoidant or resistant attachment) is a risk factor for psychopathology, and secure attachment is a protective factor particularly within high-risk groups.²¹⁻²⁵ Stronger links with psychopathology are evident for young children who exhibit disorganized attachments to their primary caregivers or other atypical attachment classifications such as insecure/other in preschool children.^{21,23,24}

The Strange Situation Procedure has been enormously useful in developmental attachment research; however, its routine clinical use is limited by the fact that it was not designed as a diagnostic procedure but rather as a laboratory observation of the balance between attachment and exploratory behaviors in young children in the presence of an attachment figure.²⁶ Classifications of attachment are not diagnoses, nor does a particular classification necessarily dictate a specific clinical approach. Some attachment-based interventions do use the Strange Situation Procedure, but

clinicians are trained in appropriate administration and interpretation in that context.²⁶

Attachment of young children to caregivers may be compromised by other risk factors that also give rise to psychiatric symptoms and disorders. This raises the question of how to define clinical disorders of attachment, that is, for those children for whom attachment disorders are the primary focus of treatment, as opposed to children with insecure or disorganized attachments who are at risk for subsequent disorders but are not currently displaying signs of a clinical disorder.

In contrast to patterns of attachment that are risk and protective factors, the *DSM-5*² and other nosologies are concerned with defining clinical disorders of attachment. In the *DSM-5*,² RAD is defined by markedly disturbed and developmentally inappropriate attachment behaviors, in which a child rarely or minimally turns preferentially to an attachment figure for comfort, support, protection, and nurturance. The child rarely seeks comfort or responds to comfort when distressed, and is socially unresponsive and has emotion regulation difficulties. Signs of the disorder appear following extremes of insufficient care.

DSED is defined in the *DSM-5*² as aberrant behavior in which a child unhesitatingly approaches and interacts with unfamiliar adults. The child shows reduced or absent reticence about approaching strangers, overly familiar verbal or physical behavior, diminished or absent checking back with an adult caregiver after venturing away, even in unfamiliar settings, and/or a willingness to “go off with” an unfamiliar adult with minimal or no hesitation. As with RAD, these behaviors follow extremes of insufficient care.

ONE DISORDER OR TWO?

Reactive attachment disorder (RAD) was divided in the *DSM-IV*²⁷ into two subtypes: an emotionally withdrawn/inhibited type and an indiscriminately social/disinhibited type. In contrast, the *International Classification of Diseases—10th Revision (ICD-10)*²⁸ defined two disorders: reactive attachment disorder, corresponding to the emotionally withdrawn/inhibited type, and disinhibited attachment disorder, corresponding to the indiscriminately social/disinhibited type. The idea was that, in the former, young children with limited opportunities to form selective attachments were withdrawn and inhibited, with no consistent displays of attachment behaviors directed to anyone. The latter, on the other hand, was intended to identify young children who similarly lacked opportunities to form selective attachments and, in response, displayed attachment behaviors indiscriminately, even to complete strangers.

Recent reviews have shown that the evidence indicates that the two different types of RAD are actually two distinct disorders.^{29–31} The two disorders arise in similar conditions of risk: experiences that limit the child’s ability to form selective attachments, such as social neglect, frequent changes in caregivers, or deprivation that may occur in institutional settings. On the other hand, they differ with regard to phenomenology, correlates, course, response to treatment, and vulnerability factors,^{29–31} and this led to them

being defined as distinct disorders in the *DSM-5*.² In this conceptualization, RAD involves disordered attachment behaviors, but in DSED, the core abnormality concerns social disinhibition. Because DSED may occur in the absence of attachment, in an aberrant attachment or in a healthy attachment to a subsequent foster or adoptive parent, one may reasonably question whether it is an attachment disorder at all.^{30,31}

CLINICAL PRESENTATION OF RAD

RAD describes a constellation of aberrant attachment behaviors and other behavioral abnormalities that are believed to result from social neglect and deprivation. For this reason, it requires a history of serious social neglect. Lack of attachment to a specific attachment figure is exceedingly rare in reasonably responsive caregiving environments. In response to severely limited opportunities to form selective attachments, young children fail to develop attachments to any caregivers—this is the essence of the disorder.^{31–33}

Children with RAD do not initiate or even show much interest in interacting with caregivers, and social reciprocity is minimal or absent. These children have limited or no positive affect and often appear unresponsive. Children with RAD do not show consistent or robustly developed attachment behaviors—in fact, they rarely seek proximity to specific adults, fail to check in with adults, even those they have been repeatedly exposed to, and neither look for nor accept comfort from caregivers in times of emotional need. They also may display episodes of unexplained irritability, sadness, or fearfulness around familiar caregivers.

COMORBIDITY OF RAD

Only limited data are available at this time about disorders that might be comorbid with RAD. Yet, given that severe childhood adversity seems to increase risk for many types of psychopathology,^{33,34} it is likely that comorbidity is the rule rather than the exception. In particular, stereotypes³⁵ and cognitive delays³⁶ are both associated with deprived caregiving environments and often co-occur with RAD. In addition, comorbidity with depressive symptoms also has been noted in young children with histories of institutional rearing.³⁷ Some maltreated children have been documented to show signs of both posttraumatic stress disorder (PTSD) and RAD,³⁸ but no studies have yet documented the degree of comorbidity between RAD and PTSD. There is, however, longitudinal evidence that disorganized attachment in infancy is associated with the development of PTSD in school-aged children who were exposed to significant traumatic events,³⁹ and children diagnosed with RAD are likely to have such trauma exposure.

DIFFERENTIAL DIAGNOSIS OF RAD

RAD must be distinguished primarily from autism spectrum disorder (ASD), global developmental delay (GDD), and depression. Children with RAD share social withdrawal and reduced social reciprocity with children with ASD. Similarly, both disorders are often associated with cognitive delays and

motor stereotypies. On the other hand, in RAD, one should not see selective impairments in symbolic representation (i.e., pretend play and language) out of proportion to cognitive level. Furthermore, the restricted interests and pre-occupations, which are part of the criteria for ASD, are not found in RAD. Children with RAD have deviant social and emotional behavior, reduced responsiveness and positive affect, and emotion regulation disturbances that are not part of global developmental delay. In depression, reduced positive affect and emotion regulation difficulties are similar to what is seen in RAD. There is no reason, however, to expect that the attachment behaviors of a young child with depression would be minimal to absent, as is the case in RAD.

COURSE OF RAD

Only one longitudinal study has examined the course of RAD.⁴⁰ The Bucharest Early Intervention Project (BEIP) was the first (and so far only) randomized clinical trial of foster care as an alternative to institutional care among young children who had been abandoned and placed in institutions. Following baseline assessments in institutional settings, half of the sample was then randomly selected for placement into a foster care network that was created by the project because foster care was largely unavailable as an option in Bucharest when the study began. The children were 6 to 30 months old at the time of initial assessment, and they were systematically re-evaluated at 30, 42, and 54 months and then at 8 years. Although BEIP included multiple brain and behavioral outcomes, RAD and DSED were always a central focus of the study because they had been identified in descriptive studies of children raised in institutions throughout the latter half of the 20th century.

BEIP results demonstrated that young children who were randomly selected to be removed from institutions and placed in foster care showed an early and substantial decrease in signs of RAD compared to children who remained institutionalized for longer periods of time.⁴¹ After placement at a mean of 22 months of age, the group in foster care had levels of RAD comparable to never-institutionalized Romanian children living with their families by 30 months of age. On the other hand, for the group randomized to remain in institutional settings, there was moderate stability in signs of RAD from infancy through age 8 years.⁴¹ Those who remained the longest in institutional care had the most persistently high signs of RAD over time.³⁷

In addition, in studies of children adopted out of institutions, there have been no cases of RAD in follow-ups conducted months to years after adoption.^{42,43} In other words, once children are removed from the socially depriving environments of institutions and are placed with families, the emotional withdrawal and inhibition characteristics of children with RAD disappear.

There is no evidence to suggest that, once removed from deprivation, children with RAD reach an age at which they no longer can form attachments, although lasting developmental compromises have been described in extreme situations of deprivation such as with feral children. Nevertheless, less is known about the risk to long-term interpersonal

relatedness and even psychopathology in individuals who have a history of RAD in early childhood. There are no data available about heterotypic continuity in RAD.

EPIDEMIOLOGY OF RAD

Few data exist about the prevalence of RAD. It is thought to be rare, but only one published community study has reported on prevalence. That study of 350 Romanian preschool children recruited from pediatric waiting rooms found no cases of RAD.⁴⁴ In fact, even in a study of young children raised in institutional settings for varying amounts of time, the rates of the emotionally withdrawn/inhibited type of RAD in 54-month-old children was only 4.1%.³⁷ On the other hand, this and another study demonstrated that as many as 40% of currently institutionalized young children show significant signs of RAD.⁸ Rates in young children newly placed in foster care are less clear, but signs of RAD have been reported in these populations.^{45,46} Beyond these extreme populations, RAD appears to be exceedingly rare. RAD is more common in children currently living in institutions and, given that RAD reliably improves when children are removed from institutional care, eliminating institutional rearing should be a priority (or improving them in contexts where they cannot be eliminated).^{47,48} Given that the *DSM-5*² criteria require a history of severely “insufficient care,” the diagnosis should be questioned in any case in which a history of social neglect cannot be documented.

ETIOLOGY AND RISK FACTORS OF RAD

In the *DSM-5*,² RAD is classified as one of a group of “trauma and stress-related disorders.” As such, the etiology of the disorder is specified in the diagnostic criteria. The phenotype of RAD must have resulted from the child’s experience of severely “insufficient caregiving,” which is operationalized as social neglect or deprivation, repeated changes of primary caregivers, or rearing in institutions with high child-to-caregiver ratios. Research has confirmed that children who have experienced adverse, neglectful caregiving environments have an increased risk of RAD compared to children who are raised in lower-risk environments.^{5,8,10,37,45,49} In fact, the phenotype of RAD has not been reported in young children in the absence of a history of neglect.

In a study of young children living in large institutions, individual differences in observed quality of caregiving were related to individual differences in signs of RAD, even after controlling for other child and environmental characteristics.¹⁰ This implies that the degree of adversity in the caregiving environment is an important factor in the appearance of RAD. Intraindividual vulnerability factors have not yet been identified that may clarify who does and does not develop RAD given the presence of known risk factors.

CLINICAL PRESENTATION OF DSED

Children with DSED also have a history of social neglect, but they are usually affectively brighter and more social than children with RAD. The essence of DSED is socially disinhibited behavior with strangers. Affected children lack

restraint around adults whom they do not know, instead approaching and engaging them. In fact, they are notably willing to leave caregivers and accompany or “go off” with complete strangers without hesitation. Children with DSED often are intrusive and lack appropriate social and physical boundaries, and are emotionally “over bright” and attention seeking. Their “friendliness” is often described by caregivers as uncomfortable, and the attention seeking can include aggressive behavior at times, although studies of aggression and indiscriminate behavior in children raised in institutions have shown that they are largely independent.^{10,37,50}

Children may exhibit signs of DSED whether or not they have formed preferred attachments. That is, some children turn preferentially for comfort to a preferred attachment figure, but they still show lack of reticence about approaching, engaging with, and even going off with strangers. Because DSED has been documented in children who have and in those who have not developed preferred attachments,^{10,12,50-53} there is reason to question why DSED should be defined as an attachment disorder. This is a major reason why the *DSM-5* has defined the disorder as disinhibited social engagement disorder, distinguishing it from RAD.²⁸

COMORBIDITY OF DSED

In addition to indiscriminate behavior, social deprivation in early childhood also produces large rates of inattention/overactivity,⁵⁴ and comorbidity of attention-deficit/hyperactivity disorder (ADHD) and DSED has been documented in preschool-³⁷ and school-aged children.⁴² Although not studied systematically, PTSD also has been reported to co-occur with DSED in young maltreated children.³⁸

Two independent groups studying children adopted out of severely deprived institutions have reported that some children display features described as “quasi-autism.”^{55,56} At age 4 years, these children met full clinical criteria for autism, but by age 6 years, they no longer showed all the features of autism. Instead, they exhibited odd relatedness, including some indiscriminate behavior, and they continued to show some peculiar interests, but they had more flexible communicative abilities than is typical in autism. Some evidence suggests that it may be social communicative deficits that are most impaired in children who experience institutional rearing.⁵⁷ When followed up farther into school age and adolescence, 7 of 15 children with quasi-autism showed indiscriminate behavior characteristic of DSED.⁵⁸ It is worth noting, however, that the majority of children with indiscriminate behavior have not been reported to show features of quasi-autism.^{7,10,37,58-63}

DIFFERENTIAL DIAGNOSIS OF DSED

Children with DSED must be distinguished from children with high levels of sociability and from children with ADHD. The distinction between highly sociable and indiscriminate behavior hinges on the degree to which the child violates accepted social norms for social boundaries and the degree of functional impairment associated with the disorder. More of each suggests DSED. For ADHD, the impulsivity is

usually behavioral and cognitive and only sometimes social, whereas in DSED, the impulsivity is social. In studies of indiscriminate behavior, cognitive inhibitory control and behavioral impulsivity, some clustering is apparent, but these do appear to be distinct phenomena.^{37,42,51,58}

COURSE OF DSED

DSED appears to follow a more variable course than RAD. The stability of indiscriminate behavior is modest to moderate, both in institutionalized³⁷ and in formerly institutionalized children.⁴² That is, longitudinal studies that have followed up children from a few years^{37,52} to more than a decade^{42,58} consistently have shown that if children raised in institutions show indiscriminate behavior when first assessed, a minority of them continue to show indiscriminate behavior, even if they are later adopted or placed with families. For example, in the BEIP, reduction in signs of DSED were less dramatic than the consistent and almost complete reduction in signs of RAD following placement in families.⁴¹

Generally, the earlier that a young child can be placed in an environment conducive to the development of preferred attachment the better, but the long-term outcomes of children diagnosed in early childhood with DSED is not well established. In children raised in institutions who were followed up from less than 30 months of age to 54 months of age, signs of DSED were predictive of overall psychiatric impairment at 54 months.³⁷ Still, little is known about individual differences in prognosis, as risk and protective factors have not been well delineated among children with this disorder. In fact, the available data suggest that there are significant individual differences in the course of DSED, with some children showing sharp and consistent declines in indiscriminate behavior and others showing long-term (years') persistence. No data are yet available about heterotypic continuity of DSED, although indiscriminate behavior in early childhood has been associated with difficulties initiating and responding competently to peer relationships in adolescents.⁴³

EPIDEMIOLOGY OF DSED

The prevalence of DSED also is not completely clear because the condition has not been subjected to large, community-based studies explicitly designed to estimate prevalence. Nevertheless, DSED occurs only in a minority of children who have been severely neglected and subsequently placed in foster care or in those raised in institutions. Even in such high-risk populations, the condition occurs in fewer than 20% of children.^{37,42,62} DSED has not been studied as thoroughly in other contexts of serious environmental risk.

Only one published study has explicitly reported on the community prevalence of DSED in preschool children. In a study of 350 Romanian preschool children recruited from pediatric clinics, Gleason *et al.*⁴⁴ reported a point prevalence for DSED of 2%. This is in contrast to the finding that even among children 2.5 to 4.5 years old, who had histories of

being raised for variable times in institutions, the rates were 17% to 18%.³⁷ Signs of indiscriminate behavior are definitely elevated in preschool children following institutional rearing^{12,37,50} and social neglect,^{45,60} although they also have been reported in a few young children with no histories of either.¹²

ETIOLOGY AND RISK FACTORS FOR DSED

DSED shares the same etiology as RAD, that is, the child with the disorder must have experienced “insufficient caregiving” severe enough to explain the phenotype. Studies of children who have been maltreated^{34,38,45,60,62} and those raised in institutions^{9,42,51,52,63} have confirmed that indiscriminate behavior, the central feature of the phenotype, is increased compared to that in children who have not experienced such extremes of care.

One group of children with high levels of indiscriminate social behavior but no evidence of social neglect are those diagnosed with Williams syndrome, which results from a deletion in the long arm of the seventh chromosome.⁶⁴ Tager-Flusberg and colleagues (personal communication, June 10, 2007) assessed a small number of children diagnosed with Williams syndrome who had no history of adverse experiences, and found that parents reported that the children had extremely high levels of indiscriminate behavior. Data on children with Williams syndrome indicate the importance of maintaining the insufficient caregiving as a criterion to define DSED to distinguish indiscriminate behavior resulting from social neglect from that due primarily to genetic and/or neurodevelopmental abnormalities. The required history of extremely insufficient caregiving also is important because otherwise it might be challenging to distinguish children with core signs of the disorder from those with high levels of sociability.

It is clear from the relative rarity of DSED that most children exposed to adverse caregiving environments do not develop the disorder, raising questions about vulnerability and protective factors. There are active explorations in this area, but results to date are only preliminary. For example, Drury *et al.*⁶⁵ showed that among children raised in institutions, those who have both the met allele of BDNF and the short allele of the serotonin transporter gene have the most signs of indiscriminate behavior while living in institutions, but also the lowest levels of indiscriminate behavior once they have been placed in the enhanced caregiving environment of high-quality foster care. Thus, specific genotype variations appear differentially susceptible to the quality of the caregiving environment. A more recent study replicated the association of indiscriminate behavior and the short allele of the serotonin transporter gene but failed to replicate the BDNF association. As the authors noted, this finding is more clearly diathesis stress than differential susceptibility.⁶⁶

EMERGING DATA IN OLDER CHILDREN

There has been considerable controversy regarding diagnosis of RAD and/or DSED in school-aged children,

adolescents or adults. This controversy relates to concern about imprecise diagnostic extension of attachment disorders to cover a variety of interpersonal difficulties in individuals who have a history of early deprivation, especially signs of oppositional defiant disorder, conduct disorder, or even psychopathy in childhood.⁶⁷ A large number of psychiatric diagnoses include children who have problems with intimate relationships, and attachment disorders were never intended to include all forms of interpersonal problems. Only when the primary presenting problem is either the social reticence core to RAD or the socially disinhibited behavior defined by DSED should those diagnoses be considered. On the other hand, there is an emerging body of research on attachment disorders in school age children and adolescents. This research has demonstrated that signs of RAD and DSED are demonstrable in school-aged children and adolescents with histories of institutional rearing and maltreatment who are adopted^{61,68-71} or placed in foster care⁷²⁻⁷⁴ but also studies identifying these disorders in impoverished populations of children in whom maltreatment is not specifically identified.^{75,76} Two studies have demonstrated discriminant validity of RAD compared to ASD⁷⁷ and ADHD.⁷⁸ These investigations, despite some variability in measures and assessment methods, represent findings from multiple research groups studying children in different countries who experienced various types of insufficient care, and have provided preliminary evidence that parents of at least some older children who have experienced insufficient care endorse signs of RAD and DSED.

Recent studies have demonstrated extensive comorbidity in older children and adolescents diagnosed with attachment disorders^{75,79} and have addressed the discriminant validity of RAD as compared to ASD.⁷⁷ In addition, intellectual disabilities, language problems, and learning difficulties all have been identified in school-aged children with RAD and DSED.⁸⁰⁻⁸⁴

A number of studies in older children and adolescents also have begun to explore the neurobiology of RAD and DSED. For example, Tottenham *et al.*⁸⁵ studied 4- to 17-year-old children with histories of institutional rearing by presenting them with images of mother versus stranger and recording amygdala activation during functional magnetic resonance imaging (fMRI) scanning. As predicted, reduced amygdala differentiation correlated with greater reports of indiscriminate behavior. In addition, later adoption (i.e., more institutional rearing) was associated with reduced amygdala discrimination and more indiscriminate behavior.

Shimada *et al.* used fMRI to assess children and adolescents with RAD and typically developing controls.⁸⁶ They reported reduced gray matter volume in the left primary visual cortex in children with RAD compared to that in typically developing children. Reduced gray matter volume was associated with an increased number of internalizing problems. Investigators suggested that the results made sense because the visual cortex is part of the neurocircuitry regulating the stress response to emotional visual images.

Reward processing has been proposed as a deficit in RAD. Two small studies from the same research group in

Japan assessed children and adolescents with RAD using functional imaging. Both studies used tasks with high and low monetary rewards. In one, investigators compared children with RAD to a group of children with ADHD and a nonaffected control group.⁷⁸ In the other, a larger group of children with RAD were compared to nonaffected controls.^{87,88} In both studies, significantly reduced activity in the striatum (caudate and nucleus accumbens) was observed during the high monetary reward condition in the children with RAD compared with other groups of children. The problem is that diminished reward sensitivity has been shown to be reduced in children who have experienced adversity,^{89,90} so whether the findings are specific to RAD is unclear.

All of these new findings notwithstanding, the major question about results from studies of RAD and DSED in older children concerns measurement. These disorders were originally defined and, until recently, studied almost exclusively in younger children. In extending criteria to older children, several different measures have been used, many of which include signs and symptoms that do not derive from criteria in established nosologies, such as avoiding eye contact and misunderstanding emotions. Even those who have used a well-developed multi-method approach, with screening, structured psychiatric interviews, and observational paradigms,^{91,92} it is unclear whether this is the same disorder that has been defined and studied in young children.

More research is needed on the continuity of RAD from early to middle childhood, because it is not completely clear whether in middle childhood RAD remains largely unchanged from early childhood, is stable but developmentally different in its phenomenology, or no longer exists as a distinct disorder. Only one study has assessed a sample longitudinally from early to middle childhood, but stability was not reported.⁴¹ A recent study using the same structured interview in early adolescence that had been used in early childhood demonstrated elevated signs of RAD in adolescence much as they had been in early childhood in children who had experienced exposure to lengthy institutionalization.⁹³

For DSED, the stability of indiscriminate behavior from early to middle childhood appears to be in the low to moderate range,^{42,52} and there has been more consistency in its definition. In adolescence, indiscriminate behavior may extend beyond relations with caregivers to peer relations, such as identifying new acquaintances as “best friends” or demonstrating shallow and frequently changing friendships.

More research on both of these disorders in older children is needed. A case series review of 100 referrals to an adoption and fostering service compared the frequency of attachment disorders identified in community referrals versus specialist referrals. Attachment disorders were identified four times more often in community referrals.⁹⁴ In addition, there was significant underidentification of more common disorders (e.g., attention, behavior, and anxiety disorders) in the community. This suggests a tendency among some clinicians to overdiagnose attachment disorders in children and adolescents with histories of maltreatment.⁹⁵

EVIDENCE BASE FOR PRACTICE PARAMETERS

In this Parameter, recommendations for best treatment practices are stated in accordance with the strength of the underlying empirical and/or clinical support, as follows:

- Clinical Standard [CS] is applied to recommendations that are based on rigorous empirical evidence (e.g., meta-analyses, systematic reviews, individual randomized controlled trials) and/or overwhelming clinical consensus
- Clinical Guideline [CG] is applied to recommendations that are based on strong empirical evidence (e.g., non-randomized controlled trials, cohort studies, case-control studies) and/or strong clinical consensus
- Clinical Option [OP] is applied to recommendations that are based on emerging empirical evidence (e.g., uncontrolled trials or case series/reports) or clinical opinion, but lack strong empirical evidence and/or strong clinical consensus
- Not Endorsed [NE] is applied to practices that are known to be ineffective or contraindicated

The strength of the empirical evidence is rated in descending order as follows:

- [rct] Randomized, controlled trial is applied to studies in which participants are randomly assigned to two or more treatment conditions
- [ct] Controlled trial is applied to studies in which participants are nonrandomly assigned to two or more treatment conditions
- [ut] Uncontrolled trial is applied to studies in which participants are assigned to one treatment condition
- [cs] Case series/report is applied to a case series or a case report

RECOMMENDATIONS

Assessment

Recommendation 1. For young children with a history of foster care, adoption, or institutional rearing, clinicians should inquire routinely about a) whether the child demonstrates attachment behaviors and b) whether the child is reticent with strangers. [CS]

The purpose of screening for RAD and DSED is to determine whether more formal assessment for these disorders is necessary. In the absence of validated screening instruments for RAD and DSED, clinicians should both ask about and observe attachment in the young child directed towards the parent/caregiver. For example, does the child turn preferentially to parent figures for comfort, support, nurturance, and protection? Does the child show stranger wariness during clinical assessments? Does the child protest separation from familiar caregivers if they leave? Second, clinicians should inquire about a history of institutional rearing, foster care, or adoption (especially international adoption). Clinicians should have a low threshold for more in-depth assessment.

To consider RAD in children older than 5 years, there should be a history of recent and severe deprivation. Once

children with RAD are placed in families, signs of the disorder seem to disappear as the child forms new attachments.^{12,37} DSED, on the other hand, does in some cases persist for years, even after the child forms attachments in families.^{37,42,52} For older children and adolescents, inquiring about indiscriminate behavior with peers is indicated. For example, does the child or adolescent claim “close” friendships with relatively new acquaintances?

Recommendation 2. The clinician conducting a diagnostic assessment of RAD and DSED should obtain direct evidence from both a history of the child’s patterns of attachment behavior with his or her primary caregivers and observations of the child interacting with these caregivers. [CS]

The AACAP Practice Parameter on assessment in infancy and early childhood includes basic approaches to clinical assessment of children under 5 years, which may be useful for evaluation of RAD.⁹⁶ The caregiver’s report of the child’s attachment behavior also can be useful. The clinician should gather a detailed history about, for example, the child’s pattern of comfort seeking, beginning with the onset of stranger wariness and progressing through to the time of assessment. In addition to comfort seeking, the clinician should inquire about separation protest, which peaks at around 18 months of age but typically continues into the preschool years.

Data about the child’s behavior in child care settings or schools may be useful as an indication of the child’s typical behavior in the absence of the parent/caregiver. Teacher reports of extreme withdrawal or indiscriminate behavior could raise suspicion about RAD or DSED.

Observational data is always helpful in the diagnosis of RAD, and asking the caregiver to separate from the child by leaving the room to elicit attachment behaviors often provides useful data.²² Observing the child’s approach to and interaction with the clinician permits an in vivo examination of the child’s behavior with strangers. Comparing the child’s behavior with familiar and unfamiliar adults is necessary for diagnosis. One observational procedure is presented in more detail below. Ideally, a complete assessment involves more than one observation, with interviews helping to determine how typical the observed behavior is.⁹⁷

Recommendation 3. The clinician may be aided in making the diagnosis of RAD and DSED by a structured observational paradigm that compares the child’s behavior with familiar and unfamiliar adults. [OP]

The caregiver–child relationship forms both the basis for assessment of signs of RAD and DSED and the nexus for monitoring interventions for RAD. Structured observations allow the clinician to capture how the child behaves with a familiar and an unfamiliar adult, especially if the interactions are conducted in parallel.⁶² A number of approaches to structuring a comprehensive assessment of a caregiver–child relationship have been described^{98,99} other than the Strange Situation Procedure, which has been extensively validated^{16,19,20,21} but is more likely to be used for research purposes. These alternative approaches generally involve some combination of episodes such as play,

teaching, and separation/reunion and involve careful observations of how the child behaves with a preferred attachment figure compared with an unfamiliar adult. Other approaches more specifically designed to assess caregiver–child attachment also have been studied.⁶²

If attached, the child should exhibit clear preferences for the attachment figure for nurturance, support, comfort, and protection. A separation is expected to be mildly stressful for young children and is often included to increase the probability of observing young children when they are motivated to seek comfort.

One possible model of assessment (to date, unvalidated) for children from 1 to 5 years of age is outlined in Table 1. The procedure described in Table 1 was designed for use by clinicians working in office or clinic settings. It can be administered without additional adults being involved, although ideally it is videotaped for later review. An observation room with a one-way mirror allows the clinician to observe the parent and child during Episode 5, but if such a setting is not available, the caregiver can later report on the child’s behavior during the clinician’s absence. The novel (scary) toy episode is included so that the clinician may observe preferential comfort seeking, but it is not essential to include. Throughout the procedure, the emphasis is on comparing the child’s behavior with the familiar attachment figure (i.e., parent/caregiver) and unfamiliar adult (i.e., clinician).

DSED can be assessed in many of the same paradigms as RAD, including the one outlined in Table 1, but it is imperative that a stranger interaction be included because behavior with unfamiliar adults comprises the crucial data necessary to make a diagnosis. Other research coding systems have been used to assess the child’s behavior with the stranger in the Strange Situation Procedure, but these are less useful clinically.^{12,52}

There is no formal scoring system for the procedure in Table 1; rather, the clinician’s conclusions about differences in the child’s behavior with familiar and unfamiliar adults are sufficient to inform the diagnostic process. Training in structured observational paradigms to assess attachment is always recommended,²⁶ although specialized centers may be necessary to obtain such training.

Recommendation 4. Clinicians should perform a comprehensive psychiatric assessment of children with RAD or DSED to determine the presence of comorbid disorders. [CS]

Research to date indicates substantial levels of psychiatric and social impairment and comorbidity in children with RAD and DSED. What evidence exists suggests that signs of RAD diminish rapidly after children are placed in good-enough caregiving environments.^{12,41,52} On the other hand, children with RAD may show continuing impairment in family and social relationships, even after signs of the disorder diminish in adequate caregiving environments.⁴¹ This suggests that careful evaluation of all aspects of the child’s functioning is necessary, even after the child is in an adequate caregiving environment.

Similarly, young children in foster care and post-institutionalized children often show high levels of indiscriminate behavior when assessed soon after entry.^{42,45}

TABLE 1 Clinical Observation of Attachment⁵⁷

Episode	Duration	Action	Observation
1	5 minutes	Clinician observes parent–child “free play.”	Note especially familiarity, comfort, and warmth in child as he/she interacts with attachment figure.
2	3 minutes	Clinician talks with, then approaches, then attempts to engage child in play.	Most young children exhibit some reticence, especially initially, about engaging with an unfamiliar adult.
3	3 minutes	Clinician picks up child and shows him/her a picture on the wall or looks out window with child.	This increases the stress for the child. Again, note the child’s comfort and familiarity with this stranger.
4	3 minutes	Caregiver picks up child and shows him/her a picture on the wall or looks out window with child.	In contrast to stranger pick-up, child should feel obviously more comfortable during this activity.
4a ^a	1 minute	Child is placed between caregiver and stranger and remote control novel (e.g., scary/exciting) toy is introduced.	Child should seek comfort preferentially from parent. If interested rather than frightened, child should share positive affect with parent.
5	3 minutes	Clinician leaves the room.	This separation should not elicit much of a reaction in the child, as the clinician is a stranger.
6	1 minute	Clinician returns.	Similarly, the child should not be much affected by the stranger’s return.
7	3 minutes	Caregiver leaves the room.	Child should definitely take notice of caregiver’s departure, although not necessarily exhibit obvious distress. If the child is distressed, the clinician should be of little comfort to the child.
8	1 minute	Caregiver returns.	Child’s reunion behavior with caregiver should be congruent with separation behavior. That is, distressed children should seek comfort, and nondistressed children should re-engage positively with caregiver, by introducing him or her to the toy or activity or talking with him or her about what occurred during the separation.

Note: The general rationale for the procedure is to compare the child’s behavior with the putative attachment figure to the child’s behavior with the stranger, especially with regard to degree of comfort, showing warmth and affection, reliance for help, cooperation, and seeking comfort when afraid or distressed.

^aOptional episode.

Clinical experience suggests that this often remits if they receive good enough care from foster or adoptive parents. Severely deprived, institutionally reared children who remain in institutions, on the other hand, are more likely to show persistent indiscriminate behavior over time.^{41,43} Research also has linked indiscriminate behavior with cognitive impairment, inattention/overactivity, externalizing problems, problems with inhibitory control, and electroencephalographic (EEG) abnormalities.^{37,51,53,54,58,61} All of these findings suggest that comprehensive assessments of children with signs of DSED are indicated, including use of structured interviews.

There is evidence that maltreated children generally do not receive adequate assessment and intervention for developmental delays, language disorders, and medical conditions.³⁴ Age-appropriate screens for developmental delays, speech and language functioning, and referral for a general pediatric examination and routine testing are often necessary. Although children with RAD or DSED may show improvements in developmental delays when placed in enhanced caregiving environments, they should receive appropriate referrals for those delays in case they need additional interventions.

Recommendation 5. The clinician should assess the safety of the current placement for previously maltreated children with negative behaviors who are at high risk for being re-traumatized. [CS]

Either an early history of social neglect, serial placements in foster care, or institutional rearing in early childhood appear to be necessary for diagnosis of RAD and DSED. For this reason, clinicians who observe signs of RAD or DSED should consider the possibility that the child has experienced neglect, unless there is a plausible alternative explanation. Children with a history of serious adversity, of course, may present with a variety of negative behaviors that are difficult for caregivers to manage. Clinical judgment regarding the appropriateness of a given placement should include consideration of family support and stability, caregiver psychopathology and response to previous interventions and willingness to take responsibility for the plight of the child, and severity and pattern of previous abuse.^{100,101} After assessment, any suspicion of previously unreported or current maltreatment requires reporting to the appropriate protective services authorities and/or law enforcement.

Treatment

Recommendation 6. The most important intervention for young children diagnosed with RAD or DSED is ensuring that they are provided with an emotionally available attachment figure. [CS]

Sensitive caregiving and psychological investment in the child—essential ingredients of healthy attachments—are the most important component of an intervention for these disorders.^{13[rct]} Although children with RAD always lack well-formed attachments to adult caregivers, children with DSED only sometimes do.^{8,10,37} That is, children with DSED may have no attachments, insecure attachments, or healthy and robust attachments to foster or adoptive parents. If they lack attachments, that should be the first priority of treatment.

The building blocks of secure attachment are interactive moments in which the caregiver's sensitively attuned behavior serves to help the child develop an internal sense of security. There are two basic psychotherapeutic modalities to help children with RAD and their caregivers attune to each other and interact more positively: working through the caregiver, and working with the caregiver-child dyad (and/or family) together. Working with the child alone will most likely occur only during assessment, because the emphasis during intervention is facilitating the construction of a lasting attachment relationship for the child with a devoted caregiver.

First, the clinician can work through the caregiver, by helping him or her learn how to establish positive interactions with a difficult-to-reach child, by helping the caregiver manage the child's behavior, or by working intensively to address the caregiver's own feelings of anxiety, frustration, or anger when needed. It is not uncommon for caregivers of children with RAD to feel disconnected from the child and to react with anger or anxiety. Patterns of discipline can become overly authoritative, leading to further disruption in the child's attachment behavior. Allowing the caregiver to tell about his or her relationship with the child and reviewing that narrative for evidence of distortion or derogation is an important part of assessment and a first step in selecting an approach to intervention. Generally, this can be done as part of the open-ended assessment of the caregiver's view of the relationship.

When a caregiver is not extremely stressed and the clinician has established, through observation and interview, that the caregiver is emotionally available and readily able to reflect on the child's feelings, it may be possible to train the caregiver as a co-therapist and work to strengthen the child's attachment with the caregiver by encouraging sensitive responsiveness. Structured intervention approaches using video review are available for those clinicians who obtain training. For example, Video-based Intervention to Promote Positive parenting (VIPP) is a brief home-based attachment intervention delivered in four home visits.¹⁰² Similarly, Circle of Security, originally developed as a group psychotherapy for parents, has recently developed a DVD-based, 8-session parenting intervention.¹⁰³ Neither of these approaches has yet been studied in children with RAD or

DSED, although they have been shown to enhance secure attachment in dyads, including those at high social risk in which children show disorganized attachment.

In cases in which caregivers are too overwhelmed for coaching techniques, other approaches should be used. It is not often possible for highly stressed caregivers who have negative perceptions of their children to maintain sensitive responsiveness until the caregivers are less stressed. Dyadic work (therapy with the child and primary caregiver together) is a second basic modality for working to address symptoms of RAD.^{104,105} Two different models of dyadic interactive therapy supported by randomized clinical trials are child-parent psychotherapy^{106[rct]} and Attachment and Biobehavioral Catch Up.^{107[rct]} Although neither has been examined formally in children with attachment disorders, each has been evaluated in children with disturbed attachment relationships.^{106[rct],107[rct],108[rct]} Child-parent psychotherapy focuses primarily on the caregiver's and child's experience of one another and on altering patterns of emotional communication in the dyad. The therapist helps the caregiver to appreciate the emotional experience of the child and its connection to the emotional experience of the caregiver. Attachment and Biobehavioral Catch-Up uses videotape review of caregiver-child interaction for the clinician to review with the caregiver while attempting to shape (mostly through suggestion and positive reinforcement) the caregiver's responses.

A basic tenet in dyadic therapy is to focus on parenting strengths as reflected in observed moments of clear caregiver-child engagement.^{99,104} Once trust is built through positive reinforcement of the caregiver, the therapist can point out and process moments of frustration and disengagement in order to begin to reshape the interactions. Reviewing videotaped sessions can be useful to enhance reflective functioning about interactive behavior that is more difficult to do in the actual moment.

Recommendation 7. For young children diagnosed with DSED, limiting contacts with noncaregiving adults may reduce signs of the disorder. [OP]

The data about effects of caregiving quality on signs of DSED are much less straightforward than the data linking caregiving quality and signs of RAD.^{10,37,45} Enhancing caregiving by placing children into foster or adoptive homes from conditions of neglect may be helpful in reducing signs of DSED, but additional measures may be indicated. Clinical experience suggests that reducing the child's exposure to persons beyond the immediate family for several months after a new placement begins may reduce or eliminate indiscriminate behavior, at least in some cases.

Recommendation 8. Clinicians should recommend adjunctive interventions for children who display aggressive and/or oppositional behavior that is comorbid with DSED. [CS]

A minority of children with DSED may have concurrent oppositional and/or aggressive behaviors, which also are sequelae of adverse experiences.³⁷ In those instances, evidence-based treatment approaches for aggression, disruptive or antisocial behavior, such as parent-child interaction

therapy for younger children^{109[rect]} or multisystemic therapy for older children,^{110[rect],111[rect]} may augment the therapeutic interventions outlined in recommendation 6.

Recommendation 9. Psychopharmacological interventions are not indicated for the core features of RAD or DSED. [NE]

No psychopharmacological intervention trials for RAD or DSED have been conducted, nor are there any indications for medication for the treatment of these disorders. However, pharmacological intervention for comorbid disorders—such as related anxiety disorders, ADHD, or mood disorders—may be indicated when comprehensive assessment demonstrates ongoing symptoms and impairment. On the other hand, the lack of available data on both short-term and long-term effects of pharmacological agents on young children's rapidly developing brains reinforces the need for a cautious approach to pharmacological intervention, particularly in preschool-aged children.¹¹²

Recommendation 10. Clinicians should not administer interventions designed to enhance attachment that involve noncontingent physical restraint or coercion (e.g., “therapeutic holding” or “compression holding”), “reworking” of trauma (e.g., “rebirthing therapy”), or promotion of regression for “reattachment” because they have no empirical support and have been associated with serious harm, including death. [NE]

Establishing authority and effective limit-setting arguably are important components of any parent–child treatment. In fact, physical restraint for extreme aggression and uncontrolled behavior is sometimes necessary for protection of the child or family members.¹¹³ However, attempting to promote “reattachment” through coerced and noncontingent holding for purposes of inducing rather than containing rage is more likely to be experienced by a child as humiliating and frightening, and these approaches should be avoided.

The risks to the child involved in these nontraditional approaches are unacceptably high. Six child deaths have been attributed to prescription of these approaches.¹¹⁴ For these reasons, the American Academy of Child and Adolescent Psychiatry, the American Psychiatric Association, and the American Professional Society on the Abuse of Children all have issued policy statements opposing coercive therapies for children with serious disturbances of attachment.^{115,116}

PARAMETER LIMITATIONS

AACAP Practice Parameters are developed to assist clinicians in psychiatric decision making. These Parameters are not intended to define the sole standard of care. As such, the Parameters should not be deemed inclusive of all proper methods of care or exclusive of other methods of care directed at obtaining the desired results. The ultimate judgment regarding the care of a particular patient must be made by the clinician in light of all of the circumstances presented by the patient and his or her family, the diagnostic and treatment options available, and available resources. &

This Parameter was developed by Charles H. Zeanah, MD, Tessa Chesher, DO, Neil Boris, MD, and the American Academy of Child and Adolescent Psychiatry (AACAP) Committee on Quality Issues (CQI): Heather J. Walter, MD, MPH, and Oscar G. Bukstein, MD, MPH, co-chairs; and Christopher Bellonci, MD, R. Scott Benson, MD, Regina Bussing, MD, Allan Chrisman, MD, John Hamilton, MD, Munya Hayek, MD, Helene Keable, MD, Carol Rockhill, MD, Matthew Siegel, MD, and Sandra Stock, MD.

The AACAP Practice Parameters are developed by the AACAP CQI in accordance with American Medical Association (AMA) policy. Parameter development is an iterative process among the primary author(s), the CQI, topic experts, and representatives from multiple constituent groups, including the AACAP membership, relevant AACAP committees, the AACAP Assembly of Regional Organizations, and the AACAP Council. Details of the Parameter development process can be accessed on the AACAP Web site. Responsibility for Parameter content and review rests with the author(s), the CQI, the CQI Consensus Group, and the AACAP Council.

AACAP develops both patient-oriented and clinician-oriented Practice Parameters. Patient-oriented Parameters provide recommendations to guide clinicians toward best assessment and treatment practices. Recommendations are based on the critical appraisal of empirical evidence (when available) and clinical consensus (when evidence is unavailable), and are graded according to the strength of the empirical and clinical support. Clinician-oriented Parameters provide clinicians with the information, stated as principles, needed to develop practice-based skills. Although empirical evidence may be available to support certain principles, principles are primarily based on clinical consensus. This Parameter is a patient-oriented Parameter.

The primary intended audience for the AACAP Practice Parameters is child and adolescent psychiatrists; however, the information contained therein may also be useful for other medical and mental health clinicians.

The authors acknowledge the following experts for their contributions to this Parameter: Mary Margaret Gleason, MD, Daniel Schechter, MD, Anna Smyke, PhD, and Edmund Sonuga-Barke, BSc, PhD.

Jennifer Medicus and Stephanie Demian served as the AACAP staff liaisons for the CQI.

This Practice Parameter was made available for review to the entire AACAP membership on the AACAP website from May 2014 to July 2014.

From February 2015 to June 2015, this Parameter was reviewed by a Consensus Group convened by the CQI. Consensus Group members and their constituent groups were as follows: Heather J. Walter, MD, MPH, co-chair; John Hamilton, MD, and Munya Hayek, MD (CQI); Karlen Lyons-Ruth, PhD, and Helen Egger, MD (topic experts); Sheila Marcus, MD (AACAP Infant and Preschool Committee); Allan Chrisman, MD (AACAP Disasters and Trauma Issues Committee); Susan Scherer, MD, and John Dunne, MD (AACAP Assembly of Regional Organizations); and Douglas Kramer, MD, and Jose Vito, MD (AACAP Council).

This Practice Parameter was approved by the AACAP Council in November 2015.

This Practice Parameter supersedes the Practice Parameter for the Assessment and Treatment of Children and Adolescents With Reactive Attachment Disorder of Infancy and Early Childhood, published in the *Journal of the American Academy of Child and Adolescent Psychiatry*, 2005;44(11):1206-1219.¹

This Practice Parameter is available on the Internet (www.aacap.org).

Disclosures: During preparation of this Parameter, Dr. Zeanah received grant support from the National Institute of Mental Health (NIMH), the Substance Abuse and Mental Health Services Administration (SAMHSA), the Norlien Foundation, and the Irving Harris Foundation. Dr. Chesher has no conflicts to disclose. Dr. Boris received research support from the Nemours Foundation. Oscar Bukstein, MD, MPH, co-chair, has intellectual property with Routledge Press. Heather Walter, MD, MPH, co-chair, has no financial relationships to disclose. Disclosures of potential conflicts of interest for all other individuals named above are provided on the AACAP Web site on the Practice Parameters page.

Correspondence to the AACAP Communications Department, 3615 Wisconsin Avenue NW, Washington, DC 20016.

0890-8567/\$36.00/©2016 Published by Elsevier Inc. on behalf of the American Academy of Child and Adolescent Psychiatry.

<http://dx.doi.org/10.1016/j.jaac.2016.08.004>

REFERENCES

- American Academy of Child and Adolescent Psychiatry. Practice parameter for the assessment and treatment of children and adolescents with reactive attachment disorder of infancy and early childhood. *J Am Acad Child Adolesc Psychiatry*. 2005;44:1206-1219.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*, 5th ed. (DSM-5). Washington, DC: American Psychiatric Association; 2013.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*, 3rd ed. (DSM-III). Washington, DC: American Psychiatric Association; 1980.
- Kraemer G. A psychobiological theory of attachment. *Behav Brain Sc*. 1992;15:493-541.
- Boris N, Aoki Y, Zeanah C. The development of infant-parent attachment: considerations for assessment. *Infants Young Child*. 1999;11:1-10.
- Bowlby J. Attachment and loss: retrospect and prospect. *Am J Orthopsychia*. 1982;52:664-678.
- Dobrova-Krol NA, Bakermans-Kranenburg MJ, Van IJzendoorn MH, Juffer F. The importance of quality of care: effects of perinatal HIV infection and early institutional rearing on preschoolers' attachment and indiscriminate friendliness. *J Child Psychol Psychiatry*. 2010;51:1368-1376.
- Smyke AT, Dumitrescu A, Zeanah CH. Attachment disturbances in young children. I: the continuum of caretaking casualty. *J Am Acad Child Adolesc Psychiatry*. 2002;41:972-982.
- Tizard B, Rees J. The effect of early institutional rearing on the behaviour problems and affectional relationships of four-year-old children. *J Child Psychol Psychiatry*. 1975;16:61-73.
- Zeanah CH, Smyke AT, Koga SF, Carlson E. Attachment in institutionalized and community children in Romania. *Child Dev*. 2005;76:1015-1028.
- Chisholm K, Carter M, Ames E, Morison S. Attachment security and indiscriminately friendly behavior in children adopted from Romanian orphanages. *Dev Psychopathol*. 1995;7:283-294.
- O'Connor T, Marvin R, Rutter M, Orlrick J, Britner P. Child-parent attachment following early institutional deprivation. *Dev Psychopathol*. 2003;15:19-38.
- Smyke AT, Zeanah CH, Fox NA, Nelson CA, Guthrie D. Placement in foster care enhances quality of attachment among young institutionalized children. *Child Dev*. 2010;81:212-223.
- Stovall KC, Dozier M. The development of attachment in new relationships: single subject analyses for 10 foster infants. *Dev Psychopathol*. 2000;12:133-156.
- Stovall-McClough KC, Dozier M. Forming attachments in foster care: infant attachment behaviors during the first 2 months of placement. *Dev Psychopathol*. 2004;16:253-271.
- Ainsworth M, Blehar M, Waters E, Wall S. *Patterns of Attachment: a Psychological Study of the Strange Situation*. Hillsdale, NJ: Erlbaum; 1978.
- van IJzendoorn MH, Sagi A. Cross-cultural patterns of attachment. In: Cassidy J, Shaver P (eds). *Handbook of Attachment: Theory, Research, and Clinical Applications*. New York: Guilford; 1999:713-735.
- Steele H, Steele M, Fonagy P. Associations among attachment classifications of mothers, fathers, and their infants. *Child Dev*. 1996;67:541-555.
- Weinfeld N, Sroufe L, Egeland B, Carlson E. Individual differences in infant-caregiver attachment: conceptual and empirical aspects of security. In: Cassidy J, Shaver P (eds). *Handbook of Attachment: Theory, Research, and Clinical Applications*. New York: Guilford Press; 2008:78-101.
- Sroufe L. The role of the infant-caregiver attachment in development. In: Belsky J, Nezworski T (eds). *Clinical Implications of Attachment*. Hillsdale, NJ: Erlbaum; 1988:18-38.
- van IJzendoorn MH, Schuengel C, Bakermans-Kranenburg MJ. Disorganized attachment in early childhood: meta-analysis of precursors, concomitants, and sequelae. *Dev Psychopathol*. 1999;11:225-250.
- Madigan S, Atkinson L, Laurin K, Benoit D. Attachment and internalizing behavior in early childhood: a meta-analysis. *Dev Psychol*. 2013;49:672.
- Fearon RP, Bakermans-Kranenburg MJ, Van IJzendoorn MH, Lapsley A, Roisman GI. The significance of insecure attachment and disorganization in the development of children's externalizing behavior: a meta-analytic study. *Child Dev*. 2010;81:435-456.
- Groh A, Roisman G, van IJzendoorn M, Bakermans-Kranenburg M, Fearon P. The significance of insecure and disorganized attachment for children's internalizing symptoms: a meta-analytic study. *Child Dev*. 2012;83:591-610.
- McGoron L, Gleason MM, Smyke AT, et al. Recovering from early deprivation: attachment mediates effects of caregiving on psychopathology. *J Am Acad Child Adolesc Psychiatry*. 2012;51:683-693.
- Zeanah CH, Berlin LJ, Boris NW. Practitioner review: clinical applications of attachment theory and research for infants and young children. *J Child Psychol Psychiatry*. 2011;52:819-833.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*, 4th ed. (DSM-IV). Washington, DC: American Psychiatric Association; 1994.
- World Health Organization. *The ICD-10 Classification of Mental and Behavioral Disorders: Clinical Descriptions and Diagnostic Guidelines*. Geneva: World Health Organization; 1992.
- Rutter M, Kreppner J, Sonuga-Barke E. Emanuel Miller Lecture: Attachment insecurity, disinhibited attachment, and attachment disorders: where do research findings leave the concepts? *J Child Psychol Psychiatry*. 2009;50:529-543.
- Zeanah C, Gleason M. Reactive attachment disorder: a review for DSM-5. Washington, DC: American Psychiatric Association; 2010.
- Zeanah CH, Gleason MM. Attachment disorders in early childhood—clinical presentation, causes, correlates, and treatment. *J Child Psychol Psychiatry*. 2015;56:207-222.
- Zeanah CH, Smyke AT. Attachment disorders and severe deprivation. In: Rutter M, Bishop D, Pine D, et al. (eds). *Rutter's Child and Adolescent Psychiatry*. London: Blackwell; 2015:795-805.
- Zeanah CH, Egger HL, Smyke AT, et al. Institutional rearing and psychiatric disorders in Romanian preschool children. *Am J Psychiatry*. 2009;166:777-785.
- Reems R. Children birth to three entering state's custody. *Infant Ment Health J*. 1999;20:166-174.
- Bos K, Fox N, Zeanah CH, Nelson CA. Stereotypes in children with a history of early institutional care. *Arch Pediatr Adolesc Med*. 2010;164:406-411.
- Nelson CA, Zeanah CH, Fox NA, Marshall PJ, Smyke AT, Guthrie D. Cognitive recovery in socially deprived young children: the Bucharest early intervention project. *Science*. 2007;318:1937-1940.
- Gleason MM, Fox NA, Drury S, et al. Validity of evidence-derived criteria for reactive attachment disorder: indiscriminately social/disinhibited and emotionally withdrawn/inhibited types. *J Am Acad Child Adolesc Psychiatry*. 2011;50:216-231.
- Hinshaw-Fuselier S, Boris N, Zeanah C. Reactive attachment disorder in maltreated twins. *Infant Ment Health J*. 1999;20:42-59.
- MacDonald HZ, Beeghly M, Grant-Knight W, et al. Longitudinal association between infant disorganized attachment and childhood post-traumatic stress symptoms. *Dev Psychopathol*. 2008;20:493-508.
- Nelson CA, Fox NA, Zeanah CH. *Romania's Abandoned Children: Deprivation, Brain Development and the Struggle for Recovery*. Cambridge, MA: Harvard University Press; 2014.
- Smyke AT, Zeanah CH, Gleason M, et al. A randomized controlled trial comparing foster care and institutional care for children with signs of reactive attachment disorder. *Am J Psychiatry*. 2012;169:508-514.
- Rutter M, Colvert E, Kreppner J, et al. Early adolescent outcomes for institutionally-deprived and non-deprived adoptees. I: disinhibited attachment. *J Child Psychol Psychiatry*. 2007;48:17-30.
- Hodges J, Tizard B. Social and family relationships of ex-institutional adolescents. *J Child Psychol Psychiatry*. 1989;30:77-97.
- Gleason MM, Zamfirescu A, Egger HL, Nelson CA, Fox NA, Zeanah CH. Epidemiology of psychiatric disorders in very young children in a Romanian pediatric setting. *Eur J Child Adolesc Psychiatry*. 2011;20:527-535.
- Oosterman M, Schuengel C. Autonomic reactivity of children to separation and reunion with foster parents. *J Am Acad Child Adolesc Psychiatry*. 2007;46:1196-1203.
- Zeanah CH, Smyke AT. Disorders of attachment. In: Zeanah CH (ed). *Handbook of Infant Mental Health*, 3rd ed. New York: Guilford Press; 2009:421-434.
- St. Petersburg-USA Orphanage Research Team. The effects of early social-emotional and relationship experience on the development of young orphanage children. The St. Petersburg-USA Orphanage Research Team. *Monogr Soc Res Child Dev*. 2008;73:vii-viii, 1-262, 294-265.
- Leiden Conference on the Development and Care of Children without Permanent Parents. The development and care of institutionally reared children. *Child Dev Perspect*. 2012;6:174-180.

49. Ayaz M, Ayaz AB, Başgöl ŞS, *et al*. 3-5 Yaş Grubu Kurum Bakımındaki Çocuklarda Ruhsal Hastalık Sıklığı ve İlişkili Etmenler. *Türk Psikiyatri Dergisi*. 2012;23:82-88.
50. Zeanah CH, Smyke AT, Dumitrescu A. Attachment disturbances in young children. II: indiscriminate behavior and institutional care. *J Am Acad Child Adolesc Psychiatry*. 2002;41:983-989.
51. Bruce J, Tarullo AR, Gunnar MR. Disinhibited social behavior among internationally adopted children. *Dev Psychopathol*. 2009;21:157-171.
52. Chisholm K. A three year follow-up of attachment and indiscriminate friendliness in children adopted from Romanian orphanages. *Child Dev*. 1998;69:1092-1106.
53. Lyons-Ruth K, Bureau JF, Riley CD, Atlas-Corbett AF. Socially indiscriminate attachment behavior in the strange situation: convergent and discriminant validity in relation to caregiving risk, later behavior problems, and attachment insecurity. *Dev Psychopathol*. 2009;21:355-372.
54. Roy P, Rutter M, Pickles A. Institutional care: associations between overactivity and lack of selectivity in social relationships. *J Child Psychol Psychiatry*. 2004;45:866-873.
55. Hoksbergen R, ter Laak J, Rijk K, van Dijkum C, Stoutjesdijk F. Post-institutional autistic syndrome in Romanian adoptees. *J Autism Dev Disord*. 2005;35:615-623.
56. Rutter M, Kreppner J, Croft C, *et al*. Early adolescent outcomes of institutionally deprived and non-deprived adoptees. III. Quasi-autism. *J Child Psychol Psychiatry*. 2007;48:1200-1207.
57. Levin AR, Fox NA, Zeanah CH, Nelson CA. Social communication difficulties and autism in previously institutionalized children. *J Am Acad Child Adolesc Psychiatry*. 2015;54:108-115.
58. Rutter M, Sonuga-Barke EJ, Castle JL. Investigating the impact of early institutional deprivation on development: background and research strategy of the English and Romanian Adoptees (ERA) study. *Monogr Soc Res Child Dev*. 2010;75:1-20.
59. Bakermans-Kranenburg M, Krol N, van IJzendoorn M. Impact of institutionalized care on attachment disorganization and insecurity of Ukrainian preschoolers: protective effect of the long variant of the serotonin transporter gene (5HTT). *Int J Behav Dev*. 2011;36:1-8.
60. Pears KC, Bruce J, Fisher PA, Kim HK. Indiscriminate friendliness in maltreated foster children. *Child Maltreat*. 2010;15:64-75.
61. Tarullo AR, Garvin MC, Gunnar MR. Atypical EEG power correlates with indiscriminately friendly behavior in internationally adopted children. *Dev Psychol*. 2011;47:417-431.
62. Boris NW, Hinshaw-Fuselier SS, Smyke AT, Scheeringa MS, Heller SS, Zeanah CH. Comparing criteria for attachment disorders: establishing reliability and validity in high-risk samples. *J Am Acad Child Adolesc Psychiatry*. 1997;43:568-577.
63. O'Connor T, Rutter M. Attachment disorder behavior following early severe deprivation: extension and longitudinal follow-up. *J Am Acad Child Adolesc Psychiatry*. 2000;39:703-712.
64. Dykens EM. Anxiety, fears, and phobias in persons with Williams syndrome. *Dev Neuropsychol*. 2003;23:291-316.
65. Drury SS, Gleason MM, Theall KP, *et al*. Genetic sensitivity to the caregiving context: the influence of 5HTTLPR and BDNF val66met on indiscriminate social behavior. *Physiol Behav*. 2012;106:728-735.
66. Mesquita AR, Belsky J, Li Z, *et al*. Institutionalization and indiscriminate social behavior: differential-susceptibility versus diathesis-stress models for the 5-HTTLPR and BDNF genotypes. *Physiol Behav*. 2015;152:85-91.
67. Levy TM, Orlans M. Attachment, trauma and healing: understanding and treating attachment disorder in children and families. Washington, DC: Child Welfare League of America; 1998.
68. Raaska H, Lapinleimu H, Sinkkonen J, *et al*. Experiences of school bullying among internationally adopted children: results from the Finnish Adoption (FINADO) study. *Child Psychiatry Hum Dev*. 2012;43:592-611.
69. Kay CL, Green JM. Social cognitive deficits and biases in maltreated adolescents in UK out-of-home care: relation to disinhibited attachment disorder and psychopathology. *Dev Psychopathol*. 2016;28:73-83.
70. Kay C, Green J, Sharma K. Disinhibited attachment disorder in UK adopted children during middle childhood: prevalence, validity and possible developmental origin. *J Abnorm Child Psychol*. 2016;44:1375-1386.
71. Miellet S, Caldara R, Gillberg C, Raju M, Minnis H. Disinhibited reactive attachment disorder symptoms impair social judgements from faces. *Psychiatry Res*. 2014;215:747-752.
72. Jonkman CS, Oosterman M, Schuengel C, Bolle EA, Boer F, Lindauer RJ. Disturbances in attachment: inhibited and disinhibited symptoms in foster children. *Child Adolesc Psychiatry Ment Health*. 2014;8:1.
73. Lehmann S, Havik OE, Havik T, Heiervang ER. Mental disorders in foster children: a study of prevalence, comorbidity and risk factors. *Child Adolesc Psychiatry Ment Health*. 2013;7:1-7.
74. Love L, Minnis H, O'Connor S. Factors associated with indiscriminate friendliness in high-risk children. *Infant Mental Health Journal*. 2015;36:427-445.
75. Pritchett R, Pritchett J, Marshall E, Davidson C, Minnis H. Reactive attachment disorder in the general population: a hidden ESSENCE disorder. *Sci World J*. 2013;818157. <http://dx.doi.org/10.1155/2013/818157>.
76. Minnis H, Macmillan S, Pritchett R, *et al*. Prevalence of reactive attachment disorder in a deprived population. *Br J Psychiatry*. 2013;202:342-346.
77. Davidson C, O'Hare A, Mactaggart F, *et al*. Social relationship difficulties in autism and reactive attachment disorder: improving diagnostic validity through structured assessment. *Res Dev Disabil*. 2015;40:63-72.
78. Mizuno K, Takiguchi S, Yamazaki M, *et al*. Impaired neural reward processing in children and adolescents with reactive attachment disorder: a pilot study. *Asian J Psychiatry*. 2015;17:89-93.
79. Elovainio M, Raaska H, Sinkkonen J, Mäkipää S, Lapinleimu H. Associations between attachment-related symptoms and later psychological problems among international adoptees: results from the FinAdo study. *Scand J Psychol*. 2015;56:53-61.
80. Kočovská E, Puckering C, Follan M, *et al*. Neurodevelopmental problems in maltreated children referred with indiscriminate friendliness. *Res Dev Disabil*. 2012;33:1560-1565.
81. Giltaij HP, Sterkenburg PS, Schuengel C. Psychiatric diagnostic screening of social maladaptive behaviour in children with mild intellectual disability: differentiating disordered attachment and pervasive developmental disorder behaviour. *J Intellect Disabil Res*. 2015;59:138-149.
82. Raaska H, Elovainio M, Sinkkonen J, Matomäki J, Mäkipää S, Lapinleimu H. Internationally adopted children in Finland: parental evaluations of symptoms of reactive attachment disorder and learning difficulties—FINADO study. *Child Care Health Dev*. 2012;38:697-705.
83. Raaska H, Elovainio M, Sinkkonen J, *et al*. Adopted children's language difficulties and their relation to symptoms of reactive attachment disorder: FinAdo study. *J Appl Dev Psychol*. 2013;34:152-160.
84. Schuengel C, Schipper JC, Sterkenburg PS, Kef S. Attachment, intellectual disabilities and mental health: research, assessment and intervention. *J Appl Res Intellect Disabil*. 2013;26:34-46.
85. Olsavsky AK, Telzer EH, Shapiro M, *et al*. Indiscriminate amygdala response to mothers and strangers after early maternal deprivation. *Biol Psychiatry*. 2013;74:853-860.
86. Shimada K, Takiguchi S, Mizushima S, *et al*. Reduced visual cortex grey matter volume in children and adolescents with reactive attachment disorder. *NeuroImage*. 2015;9:13-19.
87. Tomoda A. Preliminary evidence for impaired brain activity of neural reward processing in children and adolescents with reactive attachment disorder. *Yakugaku Zasshi*. 2015;136:711-714.
88. Takiguchi S, Fujisawa TX, Mizushima S, *et al*. Ventral striatum dysfunction in children and adolescents with reactive attachment disorder: functional MRI study. *Br J Psychiatry Open*. 2015;121-128.
89. Mehta MA, Gore-Langton E, Golemb N, Colvert E, Williams SC, Sonuga-Barke E. Hyporesponsive reward anticipation in the basal ganglia following severe institutional deprivation early in life. *J Cogn Neurosci*. 2010;22:2316-2325.
90. Hanson JL, Hariri AR, Williamson DE. Blunted ventral striatum development in adolescence reflects emotional neglect and predicts depressive symptoms. *Biol Psychiatry*. 2015;78:598-605.
91. Pritchett R, Rochat TJ, Tomlinson M, Minnis H. Risk factors for vulnerable youth in urban townships in South Africa: the potential contribution of reactive attachment disorder. *Vulnerable Child Youth Stud*. 2013;8:310-320.
92. Vervoort E, De Schipper JC, Bosmans G, Verschueren K. Screening symptoms of reactive attachment disorder: evidence for measurement invariance and convergent validity. *Int J Methods Psychiatr Res*. 2013;22:256-265.
93. Humphreys KL, Nelson CA, Fox NA, Zeanah CH. Signs of reactive attachment disorder and disinhibited social engagement disorder at age 12 years: Effects of institutional care history and high-quality foster care. *Dev Psychopathol*. (In press).
94. Woolgar M, Baldock E. Attachment disorders versus more common problems in looked after and adopted children: comparing

- community and expert assessments. *Child Adolesc Ment Health*. 2015; 20:34-40.
95. Woolgar M, Scott S. The negative consequences of over-diagnosing attachment disorders in adopted children: The importance of comprehensive formulations. *Clin Child Psychol Psychiatry*. 2014;19:355-366.
 96. American Academy of Child and Adolescent Psychiatry. Practice parameters for the assessment of infants and toddlers (0-36 months). *J Am Acad Child Adolesc Psychiatry*. 2007;44:1206-1219.
 97. Dejong M. Some reflections on the use of psychiatric diagnosis in the looked after or "in care" child population. *Clin Child Psychol Psychiatry*. 2010;15:589-599.
 98. Boris NW, Fueyo M, Zeanah CH. The clinical assessment of attachment in children under five. *J Am Acad Child Adolesc Psychiatry*. 1997;36:291-293.
 99. Powell B, Cooper G, Hoffman K, Marvin RS. *The Circle of Security Intervention*. New York: Guilford Press; 2014.
 100. Britner PA, Mossler DG. Professionals' decision-making about out-of-home placements following instances of child abuse. *Child Abuse Neglect*. 2002;26:317-332.
 101. Schechter DS, Willheim E. Disturbances of attachment and parental psychopathology in early childhood. *Child Adolesc Psychiatric Clin N Am*. 2009;18:665-686.
 102. Juffer F, Bakermans-Kranenburg MJ, van IJzendoorn MH. *Promoting Positive Parenting: an Attachment-Based Intervention*. New York: Taylor and Francis Group; 2008.
 103. Cooper G, Hoffman B, Powell B. *Circle of Security Parenting: a relationship based parenting program [DVD]*. 2009. Available at: <http://circleofsecurity.net>. Accessed November 18, 2013.
 104. Lieberman A, Zeanah C. Contributions of attachment theory to infant-parent psychotherapy and other interventions with infants and young children. In: Cassidy J, Shaver P (eds). *Handbook of Attachment*. New York: Guilford Press; 1999:555-574.
 105. Lieberman A, Van Horn P. *Psychotherapy of Infants and Young Children: Repairing the Effects of Stress and Trauma on Early Attachment*. New York: Guilford Press; 2008.
 106. Dozier M, Peloso E, Lewis E, Laurenceau JP, Levine S. Effects of an attachment-based intervention on the cortisol production of infants and toddlers in foster care. *Dev Psychopathol*. 2008;20:845-859.
 107. Lieberman AF, Weston DR, Pawl JH. Preventive intervention and outcome with anxiously attached dyads. *Child Dev*. 1991;62:199-209.
 108. Toth SL, Maughan A, Manly JT, Spagnola M, Cicchetti D. The relative efficacy of two interventions in altering maltreated preschool children's representational models: implications for attachment theory. *Dev Psychopathol*. 2002;14:877-908.
 109. Eyberg S, Nelson M, Boggs S. Evidence-based psychosocial treatments for children and adolescents with disruptive behavior. *J Clin Child Adolesc Psychol*. 2008;37:215-237.
 110. Henggeler SW, Melton GB, Smith LA, Schoenwald SK, Hanley JH. Family preservation using multisystemic treatment: long-term follow-up to a clinical trial with serious juvenile offenders. *J Child Fam Stud*. 1993;2:283-293.
 111. Ogden T, Hagen KA. Multisystemic therapy of serious behaviour problems in youth: sustainability of therapy effectiveness two years after intake. *J Child Adolesc Ment Health*. 2006;11:142-149.
 112. Gleason MM, Egger HL, Emslie GJ, *et al*. Psychopharmacological treatment for very young children: contexts and guidelines. *J Am Acad Child Adolesc Psychiatry*. 2007;46:1532-1572.
 113. American Academy of Child and Adolescent Psychiatry. Practice parameters for the prevention and management of aggressive behavior with special reference to seclusion and restraint. *J Am Acad Child Adolesc Psychiatry*. 2002;41(Suppl 41):4S-25S.
 114. Chaffin M, Hanson R, Saunders BE, *et al*. Report of the APSAC task force on attachment therapy, reactive attachment disorder, and attachment problems. *Child Maltreat*. 2006;11:76-89.
 115. American Academy of Child and Adolescent Psychiatry. Policy statement on coercive interventions for reactive attachment disorder. Approved by Council November, 2003. Reviewed May, 2008. Washington, DC: American Academy of Child and Adolescent Psychiatry.
 116. American Psychiatric Association. Policy statement on reactive attachment disorder. Washington, DC: American Psychiatric Association; 2002.